Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Easter Standard Time (EST).

From: 7/2/21 12:00 am *To*: 7/2/21 11:59 pm

Offsite Monitors

H ₂ S	Limit Reached?	Detected ^a 0 – 0 ppb	0.00 ppb	N.4	70 1
H ₂ S	No	0 – 0 ppb	0.00 ppb	N.4	7 0 1
H ₂ S	No	0 - 0 ppb	0.00 nph	NT 4	T O 1
			0.00 pp0	Not available	70 ppb
H_2S	No	0-0 ppb	0.00 ppb	Not available	70 ppb
H_2S	No	0 – 5 ppb	0.08 ppb	Not available	70 ppb
H_2S	No	0-0 ppb	0.00 ppb	Not available	70 ppb
				•	
H_2S	No	0-0 ppb	0.00 ppb	Not available	70 ppb
]	H_2S H_2S	H ₂ S No	H ₂ S No $0-0$ ppb	H ₂ S No $0 - 0$ ppb 0.00 ppb H ₂ S No $0 - 0$ ppb 0.00 ppb	H2SNo $0-0$ ppb 0.00 ppbNot availableH2SNo $0-0$ ppb 0.00 ppbNot available

^a Based on 10-minute averages.

Onsite Fenceline Monitors


Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^b	24-hr Average ^b	7-day Average	30-min AEGL
Station 1						
TAPI Analyzer	H ₂ S	No	1.0 – 102.8 ppb	14.9 ppb	6.2 ppb	600 ppb
Station 2						
TAPI Analyzer	H ₂ S	No	0.2 – 20.1 ppb	2.1 ppb	0.9 ppb	600 ppb
Station 3						
TAPI Analyzer	H_2S	No	0.2 – 21.2 ppb	6.9 ppb	3.6 ppb	600 ppb
hD 1 20	· .					


^b Based on 30-minute averages.

Notes:

ATSDR MRL	Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)
AEGL	EPA Acute Exposure Guidelines Levels
H_2S	Hydrogen Sulfide
TAPI	Teledyne API H ₂ S Analyzer
hr	Hour
min	Minute
ppb	Parts per billion
MRL Limit	Limit defined as a 14-day average value.

Wind rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.





Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors


Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.


The five stand-alone H₂S monitoring stations correlate with five of EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.


The prevailing wind direction for this report period was generally from the south to southsouthwest. Winds were between 4 and 6 miles per hour throughout the day before dropping to 2 to 4 miles per hour during the evening hours.


See wind rose diagram with aerial map figure for full wind data during this reporting period.


The following locations did not detect hydrogen sulfide above 1 part per billion (ppb) during this reporting period: Catawba Headstart, Treetops, River Chase and Millstone Creek.












Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors


Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

The prevailing wind direction for this report period was generally from the south to southsouthwest. Winds were between 4 and 6 miles per hour throughout the day before dropping to 2 to 4 miles per hour during the evening hours.

See wind rose diagram with aerial map figure for full wind data during this reporting period.







10:00

12:00

14:00

16:00

18:00

20:00

22:00

200

100

0 2. Jul

02:00

04:00

06:00

08:00