Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Easter Standard Time (EST).

From: 7/16/21 12:00 am *To:* 7/16/21 11:59 pm

Offsite Monitors

day eached?Ran Detec00-9	ted ^a		MRL
o 0-9	ppb 0 ppb		-
o 0-9	ppb 0 ppb	0 1	
	rrs oppo	0 ppb	70 ppb
	·		
o 0-12	ppb 1 ppb	1 ppb	70 ppb
	· · · · ·		
o 0 – 7	ppb 0 ppb	0 ppb	70 ppb
	·		
o 0-41	opb 0 ppb	1 ppb	70 ppb
• •	·		
o 0-0	ppb 0 ppb	0 ppb	70 ppb
[o 0-41	Io 0 -4 ppb 0 ppb	Io 0 -4 ppb 0 ppb 1 ppb

^a Based on 10-minute averages.

Onsite Fenceline Monitors


Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^b	24-hr Average ^b	7-day Average	30-min AEGL
Station 1						
TAPI Analyzer	H_2S	No	1 – 2 ppb	1 ppb	2 ppb	600 ppb
Station 2						
TAPI Analyzer	H_2S	No	0 – 19 ppb	2 ppb	3 ppb	600 ppb
Station 3						
TAPI Analyzer	H_2S	No	0-137 ppb	22 ppb	18 ppb	600 ppb


^b Based on 30-minute averages.

Notes:

ATSDR MRL	Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)
AEGL	EPA Acute Exposure Guidelines Levels
H_2S	Hydrogen Sulfide
TAPI	Teledyne API H ₂ S Analyzer
hr	Hour
min	Minute
ppb	Parts per billion
MRL Limit	Limit defined as a 14-day average value.

Wind rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.





Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors


Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.


The five stand-alone H₂S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.


Winds were 2 to 4 miles per hour in the morning hours and increased to 4 to 8 miles per hour in the afternoon. Wind came from the south to southwest the whole day.


See wind rose diagram with aerial map figure for full wind data during this reporting period.


The following site did not register H₂S concentration above 1 ppb- Millstone Creek.











Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors

Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds were 2 to 4 miles per hour in the morning hours and increased to 4 to 8 miles per hour in the afternoon. Wind came from the south to southwest the whole day.

See wind rose diagram with aerial map figure for full wind data during this reporting period.

