Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Easter Standard Time (EST).

From: 7/30/21 12:00 am To: 7/30/21 11:59 pm

Offsite Monitors

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	ATSDR 14-day MRL				
Catawba Headstart										
Acrulog PPB	H ₂ S	No	0 – 4 ppb	0.07 ppb	3.03 ppb	70 ppb				
Treetops										
Acrulog PPB	H ₂ S	No	0 – 44 ppb	3.49 ppb	2.11 ppb	70 ppb				
Liberty Hill										
Acrulog PPB	H_2S	No	0-6 ppb	0.47 ppb	0.62 ppb	70 ppb				
Riverchase Estates										
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	2.03 ppb	70 ppb				
Millstone Creek										
Acrulog PPB	H_2S	No	0-20 ppb	1.79 ppb	0.30 ppb	70 ppb				

^a Based on 10-minute averages.

Onsite Fenceline Monitors

Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^b	24-hr Average ^b	7-day Average	30-min AEGL				
Station 1										
TAPI Analyzer	H ₂ S	No	1 – 3 ppb	1.45 ppb	9.23 ppb	600 ppb				
Station 2										
TAPI Analyzer	H ₂ S	No	0 – 56 ppb	6.88 ppb	5.94 ppb	600 ppb				
Station 3										
TAPI Analyzer	H ₂ S	No	0 - 188 ppb	27.42 ppb	14.83 ppb	600 ppb				

^b Based on 30-minute averages.

Notes:

ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)

AEGL EPA Acute Exposure Guidelines Levels


H₂S Hydrogen Sulfide


TAPI Teledyne API H₂S Analyzer

hr Hour min Minute ppb Parts per billion

MRL Limit Limit defined as a 14-day average value.

Wind rose - Shows the direction the wind is coming from, the monitoring station being at the center of the rose.





Legend



Offsite Fixed Monitoring Locations



Onsite Fixed Monitoring Locations



New-Indy Catawba

Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors

Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.

The five stand-alone H₂S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.


Winds came from the south to southwest at 1 to 7 miles per hour throughout the day.

See wind rose diagram with aerial map figure for full wind data during this reporting period.


Catawba Headstart - 7/30/2021 **– – –** 70 ppb H2S Conc. (ppb) 80 70 H2S Concentration (ppb) 60 50 40 30 20 10 0 10:00 12:00 14:00 16:00 18:00 20:00 22:00 0:00 4:00 6:00 2:00 8:00

Time (EST)


Treetops - 7/30/2021




Liberty Hill - 7/30/2021



Riverchase Estates - 7/30/2021



Millstone Creek - 7/30/2021





Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors

Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds came from the south to southwest at 1 to 7 miles per hour throughout the day.





Station3 - 7/30/2021

