# **Air Monitoring Summary Tables**

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Easter Standard Time (EST).

From: 09/20/21 12:00 am To: 09/20/21 11:59 pm

Offsite Monitors

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected <sup>a</sup>	24-hr Average <sup>a</sup>	7-day Average	ATSDR 14-day MRL				
Catawba Headstart										
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.01 ppb	70 ppb				
Treetops										
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.01 ppb	70 ppb				
Liberty Hill										
Acrulog PPB	$H_2S$	No	0 – 1 ppb	0.06 ppb	0.15 ppb	70 ppb				
Riverchase Estates										
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.02 ppb	70 ppb				
Millstone Creek	Millstone Creek									
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb				

<sup>&</sup>lt;sup>a</sup> Based on 30-minute averages.

#### **Onsite Fenceline Monitors**

Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected <sup>b</sup>	24-hr Average <sup>b</sup>	7-day Average	30-min AEGL
Station 1						
TAPI Analyzer	H2S	No	1 – 46 ppb	5.80 ppb	5.20 ppb	600 ppb
Station 2						
TAPI Analyzer	H2S	No	0-0 ppb	0.20 ppb	0.55 ppb	600 ppb
Station 3						
TAPI Analyzer	H2S	No	0-0 ppb	0.20 ppb	0.62 ppb	600 ppb

<sup>&</sup>lt;sup>b</sup> Based on 30-minute averages.

### Notes:

ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)

AEGL EPA Acute Exposure Guidelines Levels


H<sub>2</sub>S Hydrogen Sulfide


TAPI Teledyne API H<sub>2</sub>S Analyzer

hr Hour min Minute ppb Parts per billion

MRL Limit Limit defined as a 14-day average value.

### Wind rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.





### Legend



Offsite Fixed Monitoring Locations



**Onsite Fixed Monitoring Locations** 



New-Indy Catawba


## Period H<sub>2</sub>S Monitoring Hydrogen Sulfide Offsite Monitors

Below are graphs for offsite locations where hydrogen sulfide (H<sub>2</sub>S) was detected during the current reporting period.


The five stand-alone H<sub>2</sub>S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.

Winds mainly came from the northeast to the east-northeast throughout the day from 2 to 8 miles per hour.


See wind rose diagram with aerial map figure for full wind data during this reporting period.




# Treetops - 9/20/2021




# Liberty Hill - 9/20/2021



# Riverchase Estates - 9/20/2021



# Millstone Creek - 9/20/2021




## Period H<sub>2</sub>S Monitoring Hydrogen Sulfide Onsite Monitors

Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H<sub>2</sub>S measured at the onsite fence line locations may not exit mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds mainly came from the northeast to the east-northeast throughout the day from 2 to 8 miles per hour.

See wind rose diagram with aerial map figure for full wind data during this reporting period.



Time (EST)

