Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Easter Standard Time (EST).

From: 12/10/21 12:00 am To: 12/10/21 11:59 pm

Offsite Monitors

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	ATSDR 14-day MRL			
Catawba Headstart									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.02 ppb	70 ppb			
Treetops									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Liberty Hill ^c									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Riverchase Estates									
Acrulog PPB	H_2S	No	0 – 0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Millstone Creek									
Acrulog PPB	H_2S	No	0 – 0 ppb	0.00 ppb	0.00 ppb	70 ppb			

^a Based on 30-minute averages.

Onsite Fenceline Monitors

Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^b	24-hr Average ^b	7-day Average	30-min AEGL				
Station 1										
TAPI Analyzer	H_2S	No	0-8 ppb	1.10 ppb	2.14 ppb	600 ppb				
Station 2										
TAPI Analyzer	H_2S	No	0 – 6 ppb	0.66 ppb	1.06 ppb	600 ppb				
Station 3										
TAPI Analyzer	H_2S	No	0 – 5 ppb	0.75 ppb	1.23 ppb	600 ppb				

^b Based on 30-minute averages.

Notes:

ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)

AEGL EPA Acute Exposure Guidelines Levels

H₂S Hydrogen Sulfide


TAPI Teledyne API H₂S Analyzer

hr Hour min Minute


ppb Parts per billion

MRL Limit Limit defined as a 14-day average value.

Wind rose - Shows the direction the wind is coming from, the monitoring station being at the center of the rose.



^c Liberty Hill back online at 14:30 EST.



Legend



Offsite Fixed Monitoring Locations



Onsite Fixed Monitoring Locations



New-Indy Catawba


Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors

Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.


The five stand-alone H₂S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.

Winds were variable through the day, generally from the south at 2 to 6 miles per hour.


See wind rose diagram with aerial map figure for full wind data during this reporting period.




Treetops - 12/10/2021




Liberty Hill - 12/10/2021



Riverchase Estates - 12/10/2021



Millstone Creek - 12/10/2021




Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors


Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds were variable through the day, generally from the south at 2 to 6 miles per hour.

See wind rose diagram with aerial map figure for full wind data during this reporting period.





Station3 - 12/10/2021 30min Avg Conc. - - - 6 - - - 600 ppb Limit 800 700 H2S Concentration (ppb) 600 500 400 300 200 100 0 10:00 12:00 14:00 6:00 8:00 16:00 18:00 20:00 22:00 0:00 2:00 4:00 0:00

Time (EST)