Air Monitoring Summary Tables

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Eastern Standard Time (EST).

From: 8/31/22 12:00 am To: 8/31/22 11:59 pm

Offsite Monitors

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	ATSDR 14-day MRL			
Catawba Headstart									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Treetops									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Liberty Hill									
Acrulog PPB	H_2S	No	0 – 3 ppb	0.15 ppb	0.05 ppb	70 ppb			
Riverchase Estates									
Acrulog PPB	H_2S	No	0 − 1 ppb	0.06 ppb	0.01 ppb	70 ppb			
Millstone Creek									
Acrulog PPB	H_2S	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			

Onsite Fenceline Monitors

Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected ^a	24-hr Average ^a	7-day Average	30-min AEGL			
Station 1									
TAPI Analyzer	H_2S	No	0 – 13 ppb	2.79 ppb ^b	1.78 ppb	600 ppb			
Station 2									
TAPI Analyzer	H_2S	No	0-3 ppb	0.62 ppb ^c	0.64 ppb	600 ppb			
Station 3									
TAPI Analyzer	H_2S	No	0-0 ppb	0.20 ppb	0.35 ppb	600 ppb			

^a Based on 30-minute averages.

Notes:

ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL)

AEGL EPA Acute Exposure Guidelines Levels

H₂S Hydrogen Sulfide

TAPI Teledyne API H₂S Analyzer


hr Hour min Minute ppb Parts per billion


MRL Limit Limit defined as a 14-day average value.


Station 1 Wind Rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.

^b The 24-hr average at this location is not represented by the full 24-hr period; a total of 5 30-minute averages are missing from the reported period due to quarterly multipoint calibration performed at Station 1.

^c The 24-hr average at this location is not represented by the full 24-hr period; a total of 6 30-minute averages are missing from the reported period due to quarterly multipoint calibration performed at Station 2.







Legend



Offsite Fixed Monitoring Locations



Onsite Fixed Monitoring Locations



New-Indy Catawba


Period H₂S Monitoring Hydrogen Sulfide Offsite Monitors

Below are graphs for offsite locations where hydrogen sulfide (H₂S) was detected during the current reporting period.

The five stand-alone H₂S monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.

Winds were mostly coming from the north-northeast at 1 to 8 mph.


See wind rose diagram with aerial map figure for full wind data during this reporting period.




Treetops - 8/31/2022

Liberty Hill - 8/31/2022


Time (EST)



Riverchase - 8/31/2022



Millstone Creek - 8/31/2022




Period H₂S Monitoring Hydrogen Sulfide Onsite Monitors


Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the H₂S measured at the onsite fence line locations may not exit the mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds were mostly coming from the north-northeast at 1 to 8 mph.

See wind rose diagram with aerial map figure for full wind data during this reporting period.





Station3 - 8/31/2022

