# **Air Monitoring Summary Tables**

The table below summarizes monitoring data collected using a portable wireless remote monitoring system. All times in Eastern Standard Time (EST).

### *From:* 10/01/22 12:00 am

10/01/22 11:59 pm

**Offsite Monitors** 

Instrument	Analyte	ATSDR MRL 14-day Avg Reached?	Concentration Range Detected <sup>a</sup>	24-hr Average <sup>a</sup>	7-day Average	ATSDR 14-day MRL			
Catawba Headstart									
Acrulog PPB	$H_2S$	No	0 - 0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Treetops									
Acrulog PPB	$H_2S$	No	0 - 0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Liberty Hill									
Acrulog PPB	$H_2S$	No	0 - 0 ppb	0.00 ppb	0.00 ppb	70 ppb			
Riverchase Estates									
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.03 ppb	70 ppb			
Millstone Creek									
Acrulog PPB	$H_2S$	No	0-0 ppb	0.00 ppb	0.00 ppb	70 ppb			

To:

**Onsite Fenceline Monitors** 


Instrument	Analyte	30-min AEGL Reached?	Concentration Range Detected <sup>a</sup>	24-hr Average <sup>a</sup>	7-day Average	30-min AEGL				
Station 1										
TAPI Analyzer	$H_2S$	No	0 – 3 ppb	0.96 ppb	1.33 ppb	600 ppb				
Station 2										
TAPI Analyzer	$H_2S$	No	1 – 6 ppb	1.63 ppb	0.85 ppb	600 ppb				
Station 3										
TAPI Analyzer	$H_2S$	No	0 - 0 ppb	0.20 ppb	0.51 ppb	600 ppb				
<sup>a</sup> Based on 30 minute averages										

<sup>a</sup> Based on 30-minute averages.

#### Notes:

ATSDR MRL Agency for Toxic Substances and Disease Registry Minimal Risk Level (MRL) EPA Acute Exposure Guidelines Levels AEGL Hydrogen Sulfide  $H_2S$ Teledyne API H<sub>2</sub>S Analyzer TAPI Hour hr min Minute ppb Parts per billion MRL Limit Limit defined as a 14-day average value.

# Station 1 Wind Rose – Shows the direction the wind is coming from, the monitoring station being at the center of the rose.






## Period H<sub>2</sub>S Monitoring Hydrogen Sulfide Offsite Monitors


Below are graphs for offsite locations where hydrogen sulfide  $(H_2S)$  was detected during the current reporting period.


The five stand-alone  $H_2S$  monitoring stations correlate with five previous EPA's Viper monitoring system which includes areas to the north-northeast and south-southwest of the New-Indy Catawba Mill.


Winds were variable throughout the day at 1 to 12 mph.

See wind rose diagram with aerial map figure for full wind data during this reporting period.









## Period H<sub>2</sub>S Monitoring Hydrogen Sulfide Onsite Monitors

Below are graphs for onsite locations during the current reporting period.

Depending on wind direction, the  $H_2S$  measured at the onsite fence line locations may not exit the mill property at reported concentrations. Wind directions from offsite locations, blowing onto mill property, will disperse ambient concentrations to lower levels prior to exiting the plant site.

Winds were variable throughout the day at 1 to 12 mph.

See wind rose diagram with aerial map figure for full wind data during this reporting period.

